

****FULL TITLE****

*ASP Conference Series, Vol. **VOLUME**, © **YEAR OF PUBLICATION***

****NAMES OF EDITORS****

Star Formation and the Interstellar Medium in Nearby Tidal Streams (SAINTS)

S. J. U. Higdon,¹ J. L. Higdon,¹ B. J. Smith,² M. Hancock,³ and C. Struck⁴

Abstract.

We compare Spitzer Infrared Spectrograph observations of SQ-A & SQ-B in Stephan’s Quintet, Ambartzumian’s knot in Arp 105, Arp 242-N3, Arp 87-N1, a bridge star forming region, NGC 5291 N and NGC 5291 S. The PAHs tend to be mainly neutral grains with a typical size of 50 - 100 carbon atoms. The interstellar radiation field is harder than typical starburst galaxies, being similar to that found in dwarf galaxies. The neon line ratios are consistent with a recent episode of star formation $\lesssim 5$ million years ago. We detect emission from $\sim 10^6 M_\odot$ of warm H_2 in SQ-A and Arp 87N1 and $\sim 10^5 M_\odot$ in SQ-B. These results are similar to those derived for the tidal dwarf galaxies (TDGs) NGC 5291 N and NGC 5291 S and are consistent with emission from photodissociation regions. Using our 8 μm images of 14 interacting systems we identify 62 tidal star forming knots (TSFKs). The estimated stellar masses range from super star cluster ($10^4 - 10^6 M_\odot$) to TDG ($\sim 10^9 M_\odot$) sizes. The median stellar mass is $10^8 M_\odot$. The stellar mass, with some scatter, scales with the 8 μm luminosity and tends to be an order of magnitude smaller than the KISS sample of star forming dwarfs. An exception to this are the TSFKs in Arp 242 which have stellar masses similar to the KISS dwarfs. The TSFKs have “notched” 3.6 - 8 μm spectral energy distributions (SEDs) characteristic of star forming regions. The TSFKs, form two distinct clumps in a mid-infrared color diagram. There are 38 red-TSFKs with $[4.5] - [8.0] > 3$ and $[3.6] - [4.5] < 0.4$. This populations has significantly enhanced non-stellar emission, most likely due to PAHs and/or hot dust, relative to normal spirals and the KISS sample of dwarfs. The second group of 21 sources has $1.2 < [4.5] - [8.0] < 3$ and $[3.6] - [4.5] < 0.4$. This population overlaps with the colors of star forming dwarf and spiral galaxies. The redder $[4.5] - [8.0]$ population tends to have the sources with a rising 8-24 μm SED while the blue population tends to contain the sources with a descending SED. The rising SED is typical of spiral and starburst galaxies with a dominant 40 – 60 K dust component and the declining SED probably indicates a dominant hot dust component.

¹ Georgia Southern University, Department of Physics, Statesboro, GA 30458

² East Tennessee State University, Department of Physics and Astronomy, Johnson City, TN 37614

³ University of California, Riverside, Riverside, CA 92521

⁴ Iowa State University, Department of Physics and Astronomy, Ames, IA 50011

1 Introduction

In addition to triggering starbursts and active galactic nuclei, mergers of dusty, gas rich disk galaxies frequently lead to the formation of tidal tails that can stretch many disk diameters from the site of the collision (Toomre 1972; Schweizer 1972; Sanders & Mirabel 1996). These structures tend to be HI rich with blue optical colors, reflecting both their origin in the outer spiral disk and on-going star formation (van der Hulst 1979; Schombert, Wallin, & Struck 1990; Mirabel, Lutz, & Maza 1991; Hibbard & van Gorkom 1996). Zwicky (1956) proposed that dwarf galaxies might form out of self-gravitating clumps within tidal tails, and indeed, concentrations of gas and star forming regions are commonly found there, ranging in size from super star clusters (SSCs, $10^4 - 10^6 M_\odot$) to tidal dwarf galaxies (TDGs, $\sim 10^9 M_\odot$).

Tidal star forming knots (TSFKs) and the formation of gravitationally bound TDGs, formed either via tidal interactions between the parent galaxies or from ram-sweeping of debris material, may play an important role in galaxy formation and evolution. Higdon et al. (2006a), hereafter HHM06, found tidal bridges and tails associated with TDGs and TSFKs in NGC 5291, indicating further tidal interaction amongst the star forming knots. TDGs may be useful as local analogs of the multiple mergers of small dwarf-like galaxies at high redshift. More importantly, dwarf galaxies are the most common galaxy type in the current epoch, and TDGs may contribute significantly to this population in some environments, for example, in compact groups (Hunsberger, Charlton, & Zaritsky 1996).

Here we present some new results from our Spitzer study of Star Formation and the Interstellar Medium in Nearby Tidal Streams (SAINTS). We have selected 12 pre-merger binary pairs with prominent optical tails and/or bridges (Smith et al. 2007; Higdon & Higdon 2008). This is complemented by our study of two more complex systems: NGC 5291 where ram-sweeping is occurring in addition to a tidal interaction, and TSFKs in Stephan's Quintet. Using the Spitzer 8 μm images we have identified 62 TSFKs. We discuss the Infrared Spectrometer (IRS, Houck et al. 2004) observations of seven bright TDG candidates along with the Infrared Array Camera (IRAC, Fazio et al. 2004) and Multiband Imaging Photometer for Spitzer (MIPS, Rieke et al. 2004) images of the full sample and address some key questions concerning the nature of TDGs and star formation in tidal streams.

2 The Nature of Tidal Dwarf Galaxies and Tidal Star Formation

Figure 1 shows the IRS Short-Low (IRS-SL) observations of NGC 5291 N and NGC 5291 S (from HHM06), SQ-A & SQ-B in Stephan's Quintet and Ambartzumian's knot in Arp 105 (Higdon & Higdon 2008), and Arp 242-N3, and Arp 87-N1, the bridge star forming region in Arp 87. The TDGs/TSFKs are rich in atomic and molecular emission features from the ISM, including fine structure lines, e.g., [Ne II] 12.81 μm , [Ne III] 15.56 μm , polycyclic aromatic hydrocarbons (PAHs) and warm H_2 . The mid-infrared spectra of dwarf galaxies are known to differ substantially from those of spirals, with weaker PAH emission features and higher [Ne III]/[Ne II] ratios (Madden et al. 2006). However, it is unknown

whether the mid-infrared spectra of most TSFKs more closely resemble those of spirals or dwarfs, since only a handful have been studied in detail with Spitzer IRS spectra to date (e.g., HHM06; Higdon & Higdon 2008). For the TDGs in NGC 5291, the spectra more closely resemble those of star forming dwarfs than spirals.

The data was analyzed using SMART (Higdon et al. 2004) and the strengths of the PAH features were measured using PAHFIT (Smith et al. 2007). In the proceeding sections the results are used to address five questions.




Figure 1. IRS-SL Spectra in the rest-frame wavelength. From top to bottom, NGC 5291 N, NGC 5291 S, Stephan's Quintet-A, Stephan's Quintet-B, Arp 87-N1, Ambartzumian's Knot in Arp 105 & Arp 242-N3.

2.1 Are the PAH Grains Large? Are they Charged ?

Draine & Li (2001) model the ratio of the 6.2 to 7.7 μm features, which are both from C-C stretching modes, to derive the PAH ion/grain size. The number of carbon atoms decreases as the ratio increases. The 11.3 μm feature is from a C-H out of plane bending mode and the ratio of the 11.3 to 7.7 μm band strength gives a measure of the PAH ion fraction. The ratio of neutral to ionized PAHs increases as the ratio increases. The PAHs in the TSFKs tend to have a strong C-H out-of-plane bending mode resulting in a mainly neutral population of PAH grains. $\text{PAH}_N \sim 80 - 90\%$ in Ambartzumian's knot, Arp 242-N3, SQ-B and Arp 87-N1 and around $\text{PAH}_N \sim 60\%$ in the remainder of the sample. Arp 242-N3 and Ambartzumian's knot in Arp 105 have small PAHs, $N_C \sim 50$. The rest of the sample have larger PAHs, $N_C \sim 100$ (the model fit to SQ-A is not well constrained).

2.2 Are the Interstellar Radiation Fields (ISRF) in TSFKs and Dwarfs Similar?

The TSFKs in SQ-A and SQ-B have $[\text{S IV}]/[\text{S III}]$ and $[\text{Ne III}]/[\text{Ne II}]$ line ratios indicating moderate excitation consistent with our earlier results for NGC 5291 N and NGC 5291 S (HHM06) and with the overlap region in the Antennae (Verma et al. 2003). The ISRF in TSFKs is harder than typical starburst

galaxies, being similar to that found in dwarfs, but softer than the ISRFs found in the extreme low metallicity Blue Compact Dwarfs (Verma et al. 2003). The neon line ratios in the TSFKs are consistent with the Starburst99 models in Thornley et al. (2000) for a recent episode of star formation $\lesssim 5$ million years ago, i.e., on-going star formation.

2.3 How Much Warm H_2 do the TSFKs Contain?

We detect emission from $\sim 10^6 M_\odot$ of warm H_2 in SQ-A and Arp 87-N1 and $\sim 10^5 M_\odot$ in SQ-B. These results are similar to those derived by HHM06 for NGC 5291 N and NGC 5291 S (HHM06). The warm H_2 masses of $10^5 - 10^6 M_\odot$ are about 100 - 1000 times smaller than the average warm molecular mass of $2 \times 10^8 M_\odot$ measured in a sample of 59 ULIRGs (Higdon et al. 2006b), but the results for both galaxy types are consistent with an origin in photodissociation regions (PDRs).




Figure 2. IRAC images of (left-right) Arp 105, Arp 242 & Stephan's Quintet. Top row $3.6 \mu\text{m}$ and bottom row $8 \mu\text{m}$. TSFKs are identified with a white circle. North is top and East is left. A square root transform is used.

2.4 Can the Mid-infrared Colors and SEDs Distinguish TSFKs from other Galaxy Types?

Figure 2 shows three examples of the $3.6 \mu\text{m}$ and $8 \mu\text{m}$ images for Arp 105, Arp 242 (The Mice) and Stephan's Quintet. The locations of the TSFKs are indicated with white circles on the figures. 95% of the sources in the 14 systems have a "notched" IRAC SED characteristic of star forming regions. 79% have $24 \mu\text{m}$ detections or meaningful upper limits. These are classified into four groups depending on the $8 - 24 \mu\text{m}$ slope. There are 18 rising ($f_{24} - f_8 > 0.3 \times f_{24}$), 12 declining SED ($f_8 - f_{24} > 0.3 \times f_{24}$), 9 flat/slowly rising ($f_{24} > f_8$ and $f_{24} - f_8 < 0.3 \times f_{24}$), and 8 flat/slowly declining ($f_{24} < f_8$ and $f_8 - f_{24} < 0.3 \times f_{24}$). The rising SED is typical of spiral and starburst galaxies with a dominant $40 - 60$ K dust component and the declining SED probably indicates a dominant hot dust component. Far-infrared observations are needed to confirm this interpretation.

Figure 3 shows that TSFKs form two distinct clumps on an IRAC 2-color diagram. The first group has 38 red-TSFKs with $[4.5] - [8.0] > 3$ and $[3.6]$




Figure 3. The horizontal and vertical lines mark the color zero-points (i.e. where the two flux densities are equal). The lower right quadrant contains sources with active star formation. The TSFKs are coded according to 8 - 24 μm SED shape: Rising - filled diamond, Flat/Rising - diamond; Descending - solid square; Flat/Descending - square; unknown - X. For comparison the Pahre et al. (2004) galaxy sample are shown as crosses and the Rosenberg et al. (2006) star forming dwarfs are shown as solid circles.

- $[4.5] < 0.4$. This population has significantly enhanced non-stellar emission, most likely due to PAHs and/or hot dust, relative to normal spirals (Pahre et al. 2004) and star forming dwarf galaxies (only overlaps with one dwarf in the KISS sample, Rosenberg et al. 2006). The second group of 21 sources has $1.2 < [4.5] - [8.0] < 3$ and $[3.6] - [4.5] < 0.4$. This population overlaps with the majority of the KISS dwarfs and the spiral galaxies from Pahre's sample. The redder $[4.5] - [8.0]$ population tends to have the sources with a rising 8-24 μm SED while the blue population tends to contain the sources with a descending SED.

2.5 What is the Range in Stellar Masses? Do the Stellar Masses Depend on the SED Type or the 8 μm Luminosity?

Preliminary stellar masses are estimated using the 4.5 μm data (e.g., Oh et al. 2008) and assuming minimal non-stellar emission in the band. We find a median $M_* = 10^8 M_\odot$, with a wide range: 4.0×10^6 to $1.3 \times 10^9 M_\odot$. The 18 rising sources have a median stellar mass of $1.0 \times 10^8 M_\odot$. The 12 declining sources have a higher median stellar mass of $4.0 \times 10^8 M_\odot$. Half of the declining sources are in Arp 242, which probably biases the median stellar mass to be greater than the average for the whole sample. The sources with flat or slowly rising/declining SEDs have stellar masses similar to that of the rising sources.

The stellar mass, with some scatter, tends to scale with the 8 μm luminosity. For a given 8 μm luminosity the KISS dwarfs have a more massive stellar component. The median stellar mass for the KISS dwarfs is $1.0 \times 10^9 M_\odot$, which is a factor of ten larger than the median value for the TSFKs. 6/8 TSFKs in

Arp 242 have declining SEDs. Four of these TSFKs overlap with 3 of the dwarfs from the KISS sample.

3 Future Work

Studies of TDGs and smaller knots of star formation in tidal features help us better understand the dwarf galaxy population as a whole. The mid-infrared emission from the ISM in TSFKs are rich in atomic, molecular and PAH emission features. The study of these systems is still in its infancy as few have been studied in detail. The next generation of telescopes, for example Herschel, JWST and ALMA offer the promise of studying both fainter objects and larger samples. In particular, such studies will provide information about what fraction are truly primordial building blocks of massive galaxies left over from an early epoch of galaxy formation, and what fraction may have been born in tidal interactions. In time we will perform a more comprehensive census of star formation and the interstellar medium in tidal streams and be able to refine models of both intergalactic enrichment and star formation triggering and regulation.

Acknowledgments. research was supported by Spitzer/NASA grants RSA No.s 1346930 (Higdon & Higdon), 1353814 (Smith) & 1347980 (Struck).

References

- Bahcall, J. N., Kirhakos, S., Saxe, D. H., & Schneider, D. P. 1997, *ApJ*, 479, 642
 Draine, B. T., & Li, A. 2001, *ApJ*, 551, 807
 Fazio, G. G., et al. 2004, *ApJS*, 154, 10
 Hibbard, J., & van Gorkom, J. 1996, *AJ*, 111, 655
 Higdon, S. J. U., et al. 2004, *PASP*, 116, 975
 Higdon, S. J. U., Higdon, J. L., & Marshall, J. 2006a, *ApJ*, 640, 768
 Higdon, S. J. U., et al. 2006b, *ApJ*, 648, 323
 Higdon, S. J. U., & Higdon, J. L. 2008, *Dark Galaxies and Lost Baryons*, Proceedings of the International Astronomical Union, IAU Symposium, 244, 356
 Houck, J., et al. 2004, *ApJS*, 154, 18
 van der Hulst, J. M., 1979, *A&A*, 71, 131
 Hunsberger, S. D., Charlton, J., & Zaritsky, D. 1996, *ApJ*, 462, 50
 Madden, S. C., et al. 2006, *Å*, 446, 877
 Mirabel, I. F., Lutz, D., & Maza, J. 1991, *A&A*, 243, 367
 Oh, S., et al. 2008, *AJ*, 136, 2761
 Pahre, M. A., et al. 2004, *ApJS*, 154, 235
 Rieke, G. H., et al. 2004, *ApJS*, 154, 25
 Rosenberg, J. L., et al. 2006, *ApJ*, 636, 742
 Sanders, D. B., & Mirabel, I. F. 1996, *ARA&A*, 34, 749
 Schombert, J. M., Wallin, J. F., & Struck, C. 1990, *ApJ*, 99, 497
 Schweizer, F., 1978, *IAU Symp. 77, Structure & Properties of Nearby Galaxies*, p. 279
 Smith, B. J., et al. 2007, *AJ*, 133, 676
 Smith, J. D. T., et al. 2007, *ApJ*, 656, 770
 Thornley, M. D., et al. 2000, *ApJ*, 539, 641
 Toomre, A., & Toomre, J. 1972, *ApJ*, 178, 623
 Verma, A., et al. 2003, *A&A*, 403, 829
 Zwicky, F., 1956, *Ergebnisse der Exakten Naturwissenschaften*, 29, 344